

Abstract

# Conceptual Model for M-Health Utilization: A Nigerian Adaptation

---

Titilayo Dorothy Odetola, B.Nursing, MSc Nursing, PhD, RN

College of Medicine, Department of Nursing, University of Ibadan, Ibadan, Nigeria

---

**Corresponding Author:**

Titilayo Dorothy Odetola, B.Nursing, MSc Nursing, PhD, RN

College of Medicine

Department of Nursing

University of Ibadan

Ibadan

Nigeria

Phone: 234 8055410093

Fax: 234 8060227398

Email: [odetolatitilayo@yahoo.com](mailto:odetolatitilayo@yahoo.com)

---

## Abstract

**Background:** Compliance with Antenatal Care (ANC) and Postnatal Care (PNC) regimen is a known determinant of pregnancy outcomes. In most developing countries, access to Skilled Birth Delivery (SBD) and Childhood Immunization (CI<sub>m</sub>), socio-cultural beliefs, physical and financial barriers significantly influence perinatal health outcome. Mobile health (M-Health) is an emerging strategy for improving healthcare utilization and compliance but the extent to which it may influence uptake of available maternal and infant welfare services in Nigeria is not documented in literature. Nurses are the fulcrum for maternal healthcare but their knowledge and roles in the use of M-Health are not known in Nigeria.

**Objective:** This study was designed to assess the effects of M-Health Nursing Intervention (MHNI) on uptake of maternal health services in Oyo South Senatorial District.

**Methods:** This quasi-experimental study involved four out of nine local government areas randomly selected and allocated into Experimental (EG) and Control (CG) groups [a semi-urban and an urban each]. All the 12 Primary Health Care facilities which had nursing personnel were purposively selected. Forty-eight nurses (EG: 21 nurses; CG: 27 nurses) and 383 literate pregnant women (EG: 191 women; CG: 192 women) at gestational age of 4-6 months, registered at the PHC were recruited consecutively. Experimental group nurses were trained on M-Health and mobile telephones were given to nurses and registered pregnant women to facilitate communication. Over an 8-month period, pregnant women received free voice calls and health promotion text messages from nurses. At baseline, 3-month and 6-month, nurses' knowledge about MHNI was assessed in EG and CG using a non-weighted 42-item pretested questionnaire. Outcome evaluation checklist was used to document utilization and completion of the following six indices among pregnant women: ANC, PNC attendance, SBD, Intermittent Preventive Treatments in pregnancy (IPT<sub>p</sub>), Tetanus Toxoid (TT), and CI<sub>m</sub> within 6 weeks of birth. Data were analyzed using descriptive statistics, Chi-square, repeated measures ANOVA and logistic regression at  $P=0.05$ .

**Results:** In the EG, knowledge score significantly increased from  $21.9 \pm 4.5$  at baseline to  $23.6 \pm 4.6$  and  $23.2 \pm 5.6$  at three-month and six-month respectively while there was no significant difference in knowledge score among CG over the study period. Comparing EG with CG, significant differences were documented in ANC attendance (66.8% versus 53.1%; OR:1.7, CI: 1.2-2.7), uptake of IPT<sub>p</sub> (47.6% versus 18.4%; OR:1.7, CI: 1.2-2.7), CI<sub>m</sub> (62.6% versus 46.9%; OR:1.9, CI: 1.3-2.8) and TT (64.5% versus 54.1%; OR:1.02, CI: 0.5-1.9), SBD (69.8% versus 36.3%; OR:1.0, CI: 0.6-1.6), PNC (69.0% versus 51.0%; OR :2.1, CI: 1.4-3.2). Significantly more women in the EG who completed ANC had IPT<sub>p</sub> (OR: 14.9, CI: 6.3-35.7); TT (OR: 8.2, CI: 1.7-39.9) and SBD (OR: 2.3, CI: 1.2-4.5) than those who did not. Likewise in CG, more women who completed ANC had IPT<sub>p</sub> (OR: 21.9, CI: 5.1-94.1) and SBD (OR: 2.0, CI: 1.1-3.8).

**Conclusions:** Mobile health nursing intervention improved uptake of maternal health services among pregnant women. Policy makers need to consider the adoption of mobile health to enhance uptake of maternal health services and improve pregnancy outcome.

(*iProc* 2015;1(1):e3) doi:[10.2196/iproc.4645](https://doi.org/10.2196/iproc.4645)

---

## KEYWORDS

antenatal care; M-health nursing intervention; postnatal care; skilled-birth delivery

(This is a conference paper presented at the Connected Health Symposium, Boston, 2015, which was not edited and is only lightly peer-reviewed).

---

## Multimedia Appendix 1

Extended abstract.

[[PDF File \(Adobe PDF File\), 349KB - iproc\\_v1i1e3\\_app1.pdf](#) ]

---

## References

---

*Edited by G Eysenbach, T Hale; submitted 11.05.15; peer-reviewed by S Odametey, E Chung; accepted 25.07.15; published 27.10.15*

*Please cite as:*

*Odetola TD*

*Conceptual Model for M-Health Utilization: A Nigerian Adaptation*

*iProc 2015;1(1):e3*

*URL: <http://www.iproc.org/2015/1/e3/>*

*doi: [10.2196/iproc.4645](https://doi.org/10.2196/iproc.4645)*

*PMID:*

©Titilayo Dorothy Odetola. Originally published in JMIR Mhealth and Uhealth (<http://www.iproc.org>), 27.10.2015. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mhealth and uhealth, is properly cited. The complete bibliographic information, a link to the original publication on <http://mhealth.jmir.org/>, as well as this copyright and license information must be included.